

Licking County Cancer Profile

Ohio Department of Health

2015



Introduction

This report provides an overview of cancer in Licking County, Ohio, including data on cancer incidence (new cases) and mortality (deaths), Ohio and U.S. comparisons, trends, stage at diagnosis, cancer screening and risk factors. This information can be used to increase awareness about the impact of cancer on Ohio residents and to develop targeted programs for cancer prevention, early detection and control.

Overview

- An average of 897 new invasive cancer cases and 339 deaths occurred each year among Licking County residents from 2008-2012.
- In 2008-2012, the cancer incidence rate for all sites/types combined in Licking County was 483.5 per 100,000, compared with the Ohio rate of 463.9 per 100,000.
- The 2008-2012 cancer mortality rate in Licking County was 189.0 per 100,000, compared with the Ohio rate of 186.6 per 100,000.
- Cancer sites/types and demographic subgroups with high numbers of cancer cases and/or high incidence rates should be prioritized in cancer prevention and early detections programs. High mortality rates may be associated with a later stage at diagnosis, lack of access to health care, inadequate treatment or other risk factors that should be addressed in cancer control initiatives.

Table 1. Average Annual Number and Age-adjusted Rates of Invasive Cancer Cases and Cancer Deaths by Sex and Race in Licking County, Ohio and the United States, 2008-2012^{1,2,3}

Demographics		Incidence				Mortality			
		Licking County Cases	Ohio Rate	U.S. Rate	Licking County Deaths	Ohio Rate	U.S. Rate		
Total		897	483.5	463.9	454.8	339	189.0	186.6	171.2
Sex	Male	456	536.0	524.2	516.6	178	233.1	227.5	207.9
	Female	441	448.1	423.1	411.2	161	160.2	158.0	145.4
Race	White	863	481.6	456.1	463.3	329	188.7	184.3	170.9
	Black	21	475.9	470.8	478.0	9	234.1	217.2	202.0


¹ Source of Ohio data: Ohio Cancer Incidence Surveillance System, Chronic Disease Epidemiology and Evaluation Section and the Bureau of Vital Statistics, Ohio Department of Health, 2015.

² Source of U.S. data: Surveillance, Epidemiology and End Results Program, National Cancer Institute and the National Center for Health Statistics, 2015.

³ Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population.

Leading Cancers


Figure 1. Percentage of Cancer Cases by Site/Type for the Leading Cancers in Licking County, 2008-2012¹



- The leading sites/types of cancer incidence in Licking County in 2008-2012 were lung and bronchus, female breast, prostate, colon and rectum, and melanoma of the skin, representing 56.0 percent of all invasive cancer cases.

¹ Source: Ohio Cancer Incidence Surveillance System, Ohio Department of Health, 2015.

Figure 2. Percentage of Cancer Deaths by Site/Type for the Leading Cancers in Licking County, 2008-2012¹



- The leading sites/types of cancer mortality in Licking County in 2008-2012 were lung and bronchus, colon and rectum, female breast, prostate and pancreas*, representing 58.9 percent of all cancer deaths.

¹ Source: Chronic Disease Epidemiology and Evaluation Section and the Bureau of Vital Statistics, Ohio Department of Health, 2015.

*Leukemia also represented 4.1 percent of cancer deaths in Licking County.

- Lung and bronchus cancer was the leading cause of cancer incidence and mortality in 2008-2012, accounting for 16.9 percent of cancer cases and 33.6 percent of cancer deaths in Licking County.

Cancer Sites/Types

Table 2. Average Annual Number and Age-adjusted Rates of Invasive Cancer Cases and Cancer Deaths by Site/Type in Licking County, Ohio and the United States, 2008-2012^{1,2,3}

Cancer Site/Type	Incidence				Mortality			
	Licking County Cases	Licking County Rate	Ohio Rate	U.S. Rate	Licking County Deaths	Licking County Rate	Ohio Rate	U.S. Rate
All Sites/Types	897	483.5	463.9	454.8	339	189.0	186.6	171.2
Bladder	44	24.1	21.9	20.3	9	5.4	5.0	4.4
Brain and Other CNS	13	7.7	6.9	6.4	9	5.1	4.5	4.3
Breast (Female)	122	123.3	120.9	124.8	23	22.9	23.6	21.9
Cervix	5	6.6	7.5	7.7	2	1.9	2.6	2.3
Colon and Rectum	79	43.4	43.1	42.4	33	18.4	17.0	15.5
Esophagus	14	7.1	5.3	4.4	10	5.6	5.0	4.2
Hodgkin Lymphoma	6	3.2	2.7	2.7	<1	*	0.4	0.4
Kidney and Renal Pelvis	37	19.2	16.4	15.6	8	4.3	4.3	3.9
Larynx	7	3.9	4.1	3.2	1	0.5	1.3	1.1
Leukemia	24	13.8	11.8	13.3	14	8.0	7.3	7.0
Liver and Intrahepatic Bile Duct	12	6.4	6.1	8.2	9	5.1	5.3	6.0
Lung and Bronchus	152	82.6	71.7	58.7	114	62.9	55.3	47.2
Melanoma of Skin	45	25.4	19.7	21.6	5	3.0	3.0	2.7
Multiple Myeloma	11	6.1	5.7	6.3	4	1.9	3.5	3.3
Non-Hodgkin Lymphoma	32	17.2	18.9	19.7	11	6.2	6.9	6.2
Oral Cavity and Pharynx	21	10.5	10.7	11.0	4	2.0	2.5	2.5
Ovary	11	10.9	11.9	12.1	9	8.4	7.9	7.7
Pancreas	16	8.8	12.4	12.4	14	7.7	11.5	10.9
Prostate	105	110.1	127.8	137.9	17	24.2	22.0	21.4
Stomach	11	5.6	6.1	7.4	3	1.6	2.9	3.4
Testis	5	6.5	5.4	5.6	<1	*	0.3	0.3
Thyroid	25	14.8	13.0	13.5	1	0.8	0.5	0.5
Uterus	32	32.0	27.7	25.1	4	4.0	4.9	4.4
Other Sites/Types	67	NA	NA	NA	35	NA	NA	NA

¹ Source of Ohio data: Ohio Cancer Incidence Surveillance System, Chronic Disease Epidemiology and Evaluation Section and the Bureau of Vital Statistics, Ohio Department of Health, 2015.

² Source of U.S. data: Surveillance, Epidemiology and End Results Program, National Cancer Institute and the National Center for Health Statistics, 2015.

³ Rates are per 100,000 and age-adjusted to the 2000 U.S. standard population. Rates are sex specific for cancers of the breast, cervix, ovary, prostate, testis and uterus.


NA = Not Applicable

CNS = Central Nervous System

* Rates may be unstable and are not presented when the count for 2008-2012 is less than five (i.e., average annual count is less than one).

Trends

Figure 3. Age-adjusted Cancer Incidence Rates in Licking County and Ohio by Year, 2003-2012^{1,2}




¹ Source: Ohio Cancer Incidence Surveillance System, Ohio Department of Health, 2015.

² Rates are per 100,000 and age-adjusted to the 2000 U.S. Standard Population.

Cancer incidence rates remained stable in Licking County and decreased 7 percent in Ohio from 2003-2012. Note: Incidence rates are often variable over time at the county level, particularly for counties with small populations.

Figure 4. Age-adjusted Cancer Mortality Rates in Licking County and Ohio by Year, 2003-2012^{1,2}



¹ Source: Chronic Disease Epidemiology and Evaluation Section and the Bureau of Vital Statistics, Ohio Department of Health, 2015.

² Rates are per 100,000 and age-adjusted to the 2000 U.S. Standard Population.

Cancer mortality rates decreased 6 percent in Licking County and 11 percent in Ohio from 2003-2012.

Stage at Diagnosis

Table 3. Percent of Cancer Cases by Stage at Diagnosis for the Leading and Screenable Cancers in Licking County, Ohio and the United States, 2008-2012^{1,2,3}

Cancer Site/Type	Licking County		Ohio		U.S.	
	Early Stage	Late Stage	Early Stage	Late Stage	Early Stage	Late Stage
Breast (Female)	69.1	28.3	67.4	29.3	70.5	27.9
Cervix	46.2	50.0	41.4	52.0	45.2	49.7
Colon and Rectum	32.9	57.1	39.9	49.7	42.5	52.1
Lung and Bronchus	16.6	70.2	16.9	69.8	18.4	75.3
Melanoma of Skin	84.3	10.5	86.1	8.6	90.6	7.2
Oral Cavity and Pharynx	20.0	72.4	29.8	64.0	45.5	50.1
Pancreas	3.7	75.6	7.5	69.4	10.5	78.4
Prostate	71.8	17.3	79.0	13.0	78.8	16.3
Testis	54.2	37.5	67.7	29.5	68.0	30.6

¹ Source of Ohio data: Ohio Cancer Incidence Surveillance System, Ohio Department of Health, 2015.

² Source of U.S. data: Surveillance, Epidemiology and End Results Program, National Cancer Institute, 2015.

³ Early stage cancers are those diagnosed at the *in situ* or local stage, and late stage cancers are those diagnosed at the regional or distant stage. Early and late stage values do not add up to 100 percent because unstaged/missing stage cancers are not included.

- Cancer stage at diagnosis is the extent or spread of the tumor from the site of origin. The stages, in order of increasing spread, are *in situ*, local, regional and distant.
- Screening programs should target cancers with high proportions of late- (regional and distant) stage tumors to increase the probability of survival.
- Cancers may be reported as unstaged or with a missing stage. High proportions of unstaged/missing stage cancers may indicate incomplete reporting and should be accounted for when comparing proportions by stage.
- Pancreatic cancer had the highest proportion of late-stage tumors in Licking County among the leading and screenable cancers.

Screening

Table 4. Percent of Adults Meeting Screening Guidelines for Female Breast, Cervical and Colon and Rectum Cancer by County Type with Comparison to Ohio, 2014^{1,2}

County Type	Female Breast*	Cervix**	Colon and Rectum***
Appalachian	72.2	76.7	61.7
Metropolitan	77.0	83.3	67.8
Rural	76.7	80.0	59.6
Suburban	72.5	81.8	65.1
Ohio	75.8	81.5	65.1

¹ Source: 2014 Ohio Behavioral Risk Factor Surveillance System, Ohio Department of Health, 2015.


² Cancer screening guidelines are based on U.S. Preventative Services Task Force recommendations.

*Female breast cancer recommendations are a mammogram every two years for women ages 50-74.

**Cervical cancer recommendations are a Pap test every three years, or a combination of Pap and HPV testing every five years, for women ages 21-65. Data shown do not include HPV co-testing and only represent women ages 21-65 who have had a Pap test in the last three years.

***Colon and rectum cancer recommendations are a screening colonoscopy every 10 years; or sigmoidoscopy every five years, with high-sensitivity fecal occult blood test (FOBT) every three years; or screening with high-sensitivity FOBT every year, for persons ages 50-74.

Figure 5. Ohio Counties by County Type, 2015¹



- There were no significant differences between county types in the percent of women meeting breast cancer screening guidelines.
- Women in metropolitan counties were most likely to meet cervical cancer screening guidelines, while women in Appalachian counties were least likely to meet the guidelines.
- Metropolitan counties had a higher percent of persons meeting colon and rectum cancer screening guidelines compared with persons residing in Appalachian and rural counties.

¹ County designations as defined by the Ohio Medicaid Assessment survey (OMAS), with the exception of Mahoning County which OMAS defines as metropolitan.

Risk Factors

Table 5. Percent of Adults ages 18+ who are Current Smokers, Do Not Meet Physical Activity Guidelines and are Overweight/Obese by County Type with Comparison to Ohio, 2013 and 2014¹

County Type	Current Smoker*	Does Not Meet Physical Activity Guidelines**	Overweight/Obese***
Appalachian	23.7	81.6	70.3
Metropolitan	21.7	80.7	67.4
Rural	21.0	82.1	65.6
Suburban	16.9	79.8	66.5
Ohio	21.0	81.0	66.7

¹ Source: 2013 and 2014 Ohio Behavioral Risk Factor Surveillance System, Ohio Department of Health, 2015. Physical activity data are from 2013, and current smoker and overweight/obesity data are from 2014.

*Current smoker is defined as persons who reported smoking at least 100 cigarettes in their lifetime and currently smoke every day or some days.

**Physical activity guidelines for adults are defined as at least 150 minutes of moderate-intensity physical activity per week and muscle-strengthening activities two or more days per week.

***Overweight classification is defined as a Body Mass Index (BMI) of 25.0-29.9 and obese classification is defined as a BMI of 30.0+.

- Adults located in suburban counties were less likely to smoke compared with those located in Appalachian and metropolitan counties.
- There were no significant differences by county type for percent of adults not meeting physical activity guidelines.
- Adults located in Appalachian counties were more likely to be overweight/obese compared with those located in rural counties.

Did You Know?

Tobacco use causes many types of cancer, including cancer of the lung, larynx (voice box), mouth, esophagus, throat, bladder, kidney, liver, stomach, pancreas, cervix, colon and rectum as well as acute myeloid leukemia.

Physical activity may reduce the risk of several types of cancer including cancer of the breast, colon and rectum, and endometrium as well as advanced prostate cancer.

If Ohioans reduce their BMI by just five percent, 23,000 cases of cancer will be prevented in Ohio by 2030, saving \$1 billion.

Sources of Data and Additional Information

Ohio Cancer Incidence Surveillance System (OCISS)

Cancer incidence data were provided by OCISS, the central cancer registry for Ohio. OCISS data can be accessed through the **Ohio Public Health Data Warehouse**, <http://publicapps.odh.ohio.gov/EDW/DataCatalog>.

Ohio Vital Statistics

Cancer mortality data were provided by the Bureau of Vital Statistics and analyzed by the Chronic Disease Epidemiology and Evaluation Section at the Ohio Department of Health.

Ohio Behavioral Risk Factor Surveillance System (BRFSS)

The BRFSS is an annual survey designed to collect data on diseases, health behaviors, clinical risk factors and injuries through landline and cell phone interviews of randomly selected adults age 18 and older. ODH conducts the Ohio BRFSS in conjunction with the Centers for Disease Control and Prevention (CDC).

U.S. Statistics

Cancer statistics for the United States were obtained from the Surveillance, Epidemiology, and End Results (SEER) Program, National Cancer Institute, and the National Center for Health Statistics, available at: <http://seer.cancer.gov>. Data sources include the *SEER Cancer Statistics Review, 1975-2012* and the SEER*Stat Database, SEER 18 Registries Research Data, released April 2015, based on the November 2014 submission. Stage distributions were calculated using SEER*Stat software version 8.2.1.

Other Sources of Information

Ohio Department of Health, Cancer Data and Statistics:
<http://www.healthy.ohio.gov/cancer/ocisshs/newrpts1.aspx>
American Cancer Society: <http://www.cancer.org>
National Cancer Institute: <http://www.cancer.gov>

The OCISS is supported in part by the State of Ohio and the CDC, National Program of Cancer Registries, cooperative agreement number 6 NU58DP003936. The contents are the sole responsibility of the authors and do not necessarily represent the official views of the CDC.

